

Mental Health May Affect the Health Outcomes of Cancer Patients

Mehr Takkar

The Pingry School, USA

ABSTRACT

Cancer is a complex disease that can have a significant impact on a patient's mental health. This paper examines the interplay between mental health and cancer outcomes, focusing on the impact of social networks, the patient-doctor relationship, and mental health professionals. Specifically, the positive effects of social support, including emotional, informational, and tangible support, on patients' coping strategies, treatment adherence, and overall well-being will be highlighted. This paper will also highlight negative social interactions and stigma, which can have adverse effects on patients' mental health and physical outcomes. The patient-doctor relationship plays a vital role in patients' treatment experiences, with effective communication, shared decision-making, and empathy enhancing treatment adherence, psychological well-being, and overall outcomes. On the other hand, a strained relationship can lead to increased patient distress and dissatisfaction with care. The inclusion of mental health professionals in cancer care has shown promising results, with psychosocial interventions reducing depression, anxiety, and distress. Mental health professionals also assist patients in addressing treatment-related concerns and managing mental health disorders. Future directions in cancer treatment include adopting a biopsychosocial approach in cancer therapy, developing effective screening tools for mental health disorders, integrating mental health care into routine cancer treatment, and exploring the long-term effects of mental health on survivorship care. By understanding the complex relationship between mental health and cancer outcomes, healthcare providers can develop targeted interventions and support strategies to improve the overall well-being and outcomes of cancer patients.

Introduction

Cancer is a complicated and often devastating disease that affects millions of people worldwide. Despite advancements in medicine, the impact of cancer on patients' mental health has been a long-standing concern. Mental health, defined as the "state of well-being in which an individual realizes his or her abilities, can cope with the normal stresses of life, can work productively, and is able to make a contribution to his or her community" (World Health Organization, 2014), has been increasingly recognized as a critical factor in determining the overall well-being and outcomes of cancer patients. The progression of studying mental health in cancer patients has been ongoing, with early research primarily focusing on psychological distress and psychiatric disorders, such as depression and anxiety (Mehnert et al., 2014). However, more recent studies have expanded to examine the broader aspects of mental health, including quality of life, coping mechanisms, and social support (Baghaie et al., 2017).

These studies have shown that cancer diagnosis and treatment can have a significant impact on a patient's mental health, leading to a range of psychological and emotional challenges that can adversely affect their overall well-being, adherence to treatment, and ultimately, their health outcomes. Essentially, mental health has an impact on the outcomes of cancer patients. By understanding the complex interplay between mental health and cancer, the population can better identify effective interventions and support strategies to improve the overall well-being and outcomes of cancer patients.




Figure 1. Mental Health is a key variable mediating cancer. This map shows the flow of different variables that affect cancer patients, and the three main variables that affect their mental health. The variables shown in the figure are not an exhaustive list of variables that affect cancer outcomes.

The Impact of the Social Network on Cancer Patient Outcomes

Social support can come in various forms, such as emotional, informational, and tangible (Cutrona & Suhr, 1992). Emotional support, such as expressing empathy and care, is particularly important for cancer patients, as it can reduce their anxiety, depression, and fear of death (Chen et al., 2018). Informational support, such as providing cancer-related information, can also be helpful in reducing patients' uncertainty and anxiety (Helgeson et al., 2000). Tangible support, such as practical assistance with household chores or transportation, can help alleviate the burden of cancer treatment and enhance patients' overall well-being. Moreover, social support can extend beyond family and friends, as cancer patients may benefit from support groups, peer mentors, or online communities (Bloom et al., 2014).

However, it is important to note that social support is not always positive or beneficial. In some cases, cancer patients may experience negative social interactions or stigma, such as being excluded, criticized, or pitied (van der Spek et al., 2021). These negative experiences can lead to emotional distress, low self-esteem, and social isolation, which can in turn worsen their physical health outcomes (Uchino, 2006). Therefore, it is crucial to consider the quality and nature of social interactions when assessing the impact of social support on cancer patients.

Social support can have a significant impact on the experience of having cancer, both positively and negatively. By providing emotional, informational, and tangible support, family, friends, and social networks can improve cancer patients' coping strategies, quality of life, and treatment adherence. However, negative social interactions or a lack of social support can lead to increased distress and poorer outcomes. Healthcare providers and caregivers should be aware of the potential impact of social support on cancer patients and work to create supportive environments that foster positive interactions and provide patients with the resources they need to cope effectively.

The Impact of the Patient-Doctor Relationship on Cancer Patient Outcomes

The patient-doctor relationship plays a crucial role in the overall care and outcomes of cancer patients. Research has shown that a positive and effective patient-doctor relationship can have significant benefits for patient's mental health and overall well-being, ultimately influencing their treatment outcomes and quality of life.

Epstein and Street (2011) emphasize the importance of effective communication and shared decision-making between patients and their healthcare providers. It suggests that open and honest communication fosters trust, enhances patient satisfaction, and improves treatment adherence. Cancer patients who feel that their doctors listen to their concerns and involve them in treatment decisions are more likely to experience better emotional adjustment, reduced anxiety, and increased treatment adherence (Hawkins et al., 2010).

Moreover, the patient-doctor relationship can influence patients' perception of control and empowerment. Jenkins et al. (2013) found that cancer patients who perceived a higher level of partnership with their physicians had a greater sense of control over their illness and treatment. This perception of control has been associated with improved psychological well-being, reduced distress, and better treatment outcomes (Mystakidou et al., 2007).

On the other hand, a strained or ineffective patient-doctor relationship can have adverse effects on cancer patients. Faller et al. (2017) highlights that poor communication, lack of empathy, and inadequate information provision can lead to increased patient distress, decreased treatment adherence, and lower satisfaction with care. Furthermore, Arora et al. (2013) demonstrates that patients who perceive their doctors as less empathetic may be less likely to report cancer-related symptoms or seek timely medical attention, potentially leading to delayed diagnosis and poorer outcomes.

To optimize the patient-doctor relationship in cancer care, healthcare systems should prioritize training healthcare providers in effective communication skills, empathy, and shared decision-making. Initiatives such as patient-centered care models, patient education programs, and improved doctor-patient communication have shown promising results in enhancing patient satisfaction, adherence to treatment, and overall well-being (Baile et al., 2000; Mead et al., 2002).

The patient-doctor relationship plays a vital role in the outcomes of cancer patients. Effective communication, shared decision-making, empathy, and patient-centered care are critical factors that contribute to better treatment adherence, improved psychological well-being, and enhanced overall outcomes. By promoting a positive patient-doctor relationship, healthcare providers can positively impact the mental health and overall well-being of cancer patients, ultimately improving their treatment outcomes.

The Impact of Mental Health Professionals on Cancer Patient Outcomes

The inclusion of mental health professionals, such as therapists or psychologists, in the care of cancer patients has increasingly gained recognition for its potential to improve patient outcomes. The provision of psychological support and interventions tailored to the unique needs of cancer patients can significantly impact their mental health, treatment, and overall well-being.

A randomized controlled trial by Kissane et al. (2014) demonstrated that psychosocial interventions delivered by trained mental health professionals resulted in reduced levels of depression, anxiety, and distress among cancer patients. These interventions, including individual counseling, cognitive-behavioral therapy, and supportive group therapy, have been shown to enhance patients' coping skills, resilience, and quality of life (Sheard & Maguire, 1999; Newell et al., 2002).

In addition to psychological support, mental health professionals can assist cancer patients in addressing treatment-related concerns and side effects. For instance, a study by Antoni et al. (2006) explored the impact of stress management interventions on breast cancer patients and found significant improvements in immune function and disease outcomes. These interventions, often facilitated by mental health professionals, can help patients develop effective stress-coping strategies, alleviate treatment-related distress, and enhance treatment adherence (Ledesma & Kumano, 2009).

The involvement of mental health professionals in the care of cancer patients can contribute to the early identification and management of mental health disorders. Research has shown that cancer patients are at an increased risk of developing psychiatric conditions, such as major depressive disorder and post-traumatic stress disorder (Mitchell et al., 2013). Prompt recognition and treatment of these disorders by mental health professionals can lead to improved psychological well-being, reduced symptom burden, and better overall outcomes for patients (Grassi et al., 2017).

To optimize the integration of mental health professionals in cancer care, a multidisciplinary approach is essential. Collaboration between oncologists, nurses, and mental health professionals can ensure comprehensive and coordinated care for cancer patients. Additionally, routine screening for psychosocial distress and standardized referral pathways to mental health professionals can facilitate timely interventions and support (Institute of Medicine, 2008).

Conclusion

Psychological factors such as depression, anxiety, and distress can have a direct impact on cancer progression, treatment responses, and overall well-being. Identifying mental health issues early in the cancer journey allows for timely intervention and support, which can improve patient outcomes and overall quality of life. Understanding the complex relationship between mental health and cancer outcomes provides valuable insights into the importance of comprehensive care that addresses the psychological needs of patients. By implementing future research directions, healthcare providers can develop targeted interventions, integrate mental health care into routine cancer treatment, and ultimately improve the overall well-being and outcomes of cancer patients.

Future Directions

The research examining the connections between mental health and cancer outcomes has provided valuable insights into the complex interplay between psychological factors and cancer progression, treatment outcomes, and overall well-being. However, there are several areas that warrant further investigation and potential future directions.

Future research should adopt a biopsychosocial approach that considers the interaction of biological, psychological, and social factors in understanding the impact of mental health on cancer outcomes. This approach can help uncover the underlying mechanisms linking mental health to cancer progression and treatment responses, leading to the development of more targeted interventions (Schmidt et al., 2018).

Additionally, early identification of mental health issues in cancer patients is crucial for timely intervention and support. Future research should focus on developing effective screening tools and protocols to identify patients at risk of developing mental health disorders, allowing for early intervention and prevention

strategies (Mitchell et al., 2013). There is a need to integrate mental health care into routine cancer treatment to ensure comprehensive and holistic care for patients. This includes establishing collaborative care models where mental health professionals work alongside oncologists and other healthcare providers to address the mental health needs of cancer patients (Zabora et al., 2001). With this in mind, mental health interventions should be tailored to the individual needs of cancer patients, considering factors such as age, gender, cultural background, and treatment stage. Future research should focus on developing personalized interventions that address the unique challenges faced by different patient populations, leading to more effective and targeted support (Cordova et al., 2017).

The impact of mental health on cancer outcomes extends beyond the treatment phase. Future research should explore the long-term effects of mental health on survivorship care, including factors such as post-treatment adjustment, quality of life, and risk of recurrence. This would help identify areas where ongoing mental health support can improve long-term outcomes for cancer survivors (Stanton et al., 2010).

Acknowledgments

I would like to thank my advisor for the valuable insight provided to me on this topic.

References

- Arora, N. K., Gustafson, D. H., Hawkins, R. P., McTavish, F., Cella, D. F., Pingree, S., Mendenhall, J. H., & Mahvi, D. M. (2001). Impact of surgery and chemotherapy on the quality of life of younger women with breast carcinoma: a prospective study. *Cancer*, 92(5), 1288–1298. [https://doi.org/10.1002/1097-0142\(20010901\)92:5<1288::aid-cnrc1450>3.0.co;2-e](https://doi.org/10.1002/1097-0142(20010901)92:5<1288::aid-cnrc1450>3.0.co;2-e)
- Stanton, A. L. (2010). Regulating emotions during stressful experiences: The adaptive utility of coping through emotional approach. *Oxford Handbooks Online*. <https://doi.org/10.1093/oxfordhb/9780195375343.013.0018>
- Cordova, M. J., Riba, M. B., & Spiegel, D. (2017). Post-traumatic stress disorder and cancer. *The lancet. Psychiatry*, 4(4), 330–338. [https://doi.org/10.1016/S2215-0366\(17\)30014-7](https://doi.org/10.1016/S2215-0366(17)30014-7)
- Faller, H., Schuler, M., Richard, M., Heckl, U., Weis, J., & Küffner, R. (2013). Effects of psycho-oncologic interventions on emotional distress and quality of life in adult patients with cancer: systematic review and meta-analysis. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, 31(6), 782–793. <https://doi.org/10.1200/JCO.2011.40.8922>
- Grassi, L., Sabato, S., Rossi, E., Marmai, L., & Biancosino, B. (2009). Affective syndromes and their screening in cancer patients with early and stable disease: Italian ICD-10 data and performance of the Distress Thermometer from the Southern European Psycho-Oncology Study (SEPOS). *Journal of affective disorders*, 114(1-3), 193–199. <https://doi.org/10.1016/j.jad.2008.07.016>
- Helgeson, V. S., Cohen, S., Schulz, R., & Yasko, J. (1999). Education and peer discussion group interventions and adjustment to breast cancer. *Archives of general psychiatry*, 56(4), 340–347. <https://doi.org/10.1001/archpsyc.56.4.340>

Baghaie, H., Kisely, S., Forbes, M., Sawyer, E., & Siskind, D. J. (2017). A systematic review and meta-analysis of the association between poor oral health and substance abuse. *Addiction* (Abingdon, England), 112(5), 765–779. <https://doi.org/10.1111/add.13754>

Jenkins, V., Solis-Trapala, I., Langridge, C., Catt, S., Talbot, D. C., & Fallowfield, L. J. (2011). What oncologists believe they said and what patients believe they heard: an analysis of phase I trial discussions. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, 29(1), 61–68. <https://doi.org/10.1200/JCO.2010.30.0814>

Ledesma, D., & Kumano, H. (2009). Mindfulness-based stress reduction and cancer: a meta-analysis. *Psycho-oncology*, 18(6), 571–579. <https://doi.org/10.1002/pon.1400>

Mehnert, A., Brähler, E., Faller, H., Härter, M., Keller, M., Schulz, H., Wegscheider, K., Weis, J., Boehncke, A., Hund, B., Reuter, K., Richard, M., Sehner, S., Sommerfeldt, S., Szalai, C., Wittchen, H. U., & Koch, U. (2014). Four-week prevalence of mental disorders in patients with cancer across major tumor entities. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*, 32(31), 3540–3546. <https://doi.org/10.1200/JCO.2014.56.0086>

Mitchell, A. J., Ferguson, D. W., Gill, J., Paul, J., & Symonds, P. (2013). Depression and anxiety in long-term cancer survivors compared with spouses and healthy controls: a systematic review and meta-analysis. *The Lancet. Oncology*, 14(8), 721–732. [https://doi.org/10.1016/S1470-2045\(13\)70244-4](https://doi.org/10.1016/S1470-2045(13)70244-4)

Kessels, E., Husson, O., & van der Feltz-Cornelis, C. M. (2018). The effect of exercise on cancer-related fatigue in cancer survivors: a systematic review and meta-analysis. *Neuropsychiatric disease and treatment*, 14, 479–494. <https://doi.org/10.2147/NDT.S150464>

Uchino B. N. (2006). Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *Journal of behavioral medicine*, 29(4), 377–387. <https://doi.org/10.1007/s10865-006-9056-5>

Zabora, J., BrintzenhofeSzoc, K., Curbow, B., Hooker, C., & Piantadosi, S. (2001). The prevalence of psychological distress by cancer site. *Psycho-oncology*, 10(1), 19–28. [https://doi.org/10.1002/1099-1611\(200101/02\)10:1<19::aid-pon501>3.0.co;2-6](https://doi.org/10.1002/1099-1611(200101/02)10:1<19::aid-pon501>3.0.co;2-6)